

The X-ray afterglow of the Gamma-ray burst of May 8, 1997: spectral variability and possible evidence of an iron line

L. Piro¹, E. Costa¹, M. Feroci¹, F. Frontera²³, L. Amati², D. Dal Fiume², L. A. Antonelli⁴, J. Heise⁵, J. in 't Zand⁵, A. Owens⁶, A.N. Parmar⁶, G. Cusumano⁷, M. Vietri⁸,
G.C. Perola⁸

Received _____; accepted _____

¹Istituto Astrofisica Spaziale, C.N.R., Via Fosso del Cavaliere 100, 00133 Rome

² Istituto TeSRE, C.N.R., Via Gobetti 101, 40129 Bologna, Italy

³Dip. Fisica, Universita' Ferrara, Via Paradiso 12, Ferrara, Italy

⁴BeppoSAX Scientific Data Center; present address: Osservatorio Astronomico di Roma

⁵ Space Research Organization in the Netherlands, Sorbonnelaan 2, 3584 CA Utrecht,
The Netherlands

⁶Space Science Department of ESA, ESTEC, 2200 AG Noordwijk, The Netherlands

⁷Istituto Fisica Cosmica e Appl. Calc. Informatico, C.N.R., Via La Malfa 153, 90138
Palermo, Italy

⁸Dipartimento di Fisica, Universita' Roma Tre, Via della Vasca Navale 84, 00146 Roma,
Italy

ABSTRACT

We report the possible detection (99.3% of statistical significance) of redshifted Fe iron line emission in the X-ray afterglow of Gamma-ray burst GRB970508 observed by BeppoSAX. Its energy is consistent with the redshift of the putative host galaxy determined from optical spectroscopy. The line disappeared ~ 1 day after the burst. We have also analyzed the spectral variability during the outburst event that characterizes the X-ray afterglow of this GRB. The spectrum gets harder during the flare, turning to steep when the flux decreases. The variability, intensity and width of the line indicate that the emitting region should have a mass $\gtrsim 0.5M_{\odot}$ (assuming the iron abundance similar to its solar value), a size of $\sim 3 \times 10^{15}cm$, is distributed anisotropically, and is moving with sub-relativistic speed. In contrast to the fairly clean environment expected in the merging of two neutron stars, the observed line properties would imply that the site of the burst is embedded in a large mass of material, consistent with pre-explosion ejecta of a very massive star. This material could be related with the outburst observed in the afterglow 1 day after the GRB and with the spectral variations measured during this phase.

Subject headings: gamma rays: bursts – X-rays: general – line: formation

1. Introduction

Distance - scale determination of Gamma-ray bursts (GRB) has been one of the most important achievements of astrophysics in recent years. Accurate and fast localization of the prompt and afterglow emission bursts (i.e. Costa *et al.* 1997) by the Italian/Dutch X-ray satellite BeppoSAX (Piro, Scarsi & Butler 1995, Boella *et al.* 1997) led to the identification of optical counterparts (van Paradijs *et al.* 1997) and ultimately to spectral measurements of a redshift (Metzger *et al.* 1997). While the extragalactic origin of GRB has gathered solid evidence in its support, the source of the large energy implied by their distance is still speculative.

Direct information on the nature of the central engine of GRB can be derived by studying the nearby environment; for example using line spectroscopy. The star forming region and massive pre-explosion winds associated with the hypernova scenario (Paczynsky 1998) imply a mass rich environment down to short ($\sim 10^{15} \text{cm}$) distance scales. In contrast, NS-NS merging should happen in a fairly clean environment, because such objects are expected to form with significant speeds which can lead them at least a few parsecs away from the regions of star formation in which they formed. Current line measurements are inconclusive, because all the spectral features observed so far are signatures produced by the host galaxy rather than at the burst site. GRB980425 is an exception, but its tantalizing association with SN 1998bw (Galama *et al.* 1998, Kulkarni *et al.* 1998) requires clarification.

The measurement of X-ray lines emitted directly by the GRB or its afterglow could provide a direct measurement of the distance and probe into the nature of the central environment (Perna & Loeb 1998). Fe K line emission is very promising in this respect. Theoretical computations have been carried out by several authors in the framework of GRB (Mészáros & Rees 1998, Boettcher *et al.* 1999, Ghisellini *et al.* 1998). Most of this

work deals with features generated within the interstellar medium (ISM) of the host galaxy, i.e. on scale larger than several parsecs. However, even in the favourable case of dense regions of stellar formation (e.g. the site of GRB formation in the hypernova scenario), iron emission lines would be hard to detect with current X-ray instrumentation. Lately, Mészáros and Rees (1998) have shown that the circumburst environment created by the stellar wind before the explosion of the hypernova could yield a line of substantial intensity. A similarly favourable situation should be expected in a variation of the hypernova scenario, – the SupraNova (Vietri & Stella 1998), where the GRB is shortly preceded by a supernova explosion with the ejection of an iron-rich massive shell. In these cases, and for a isotropic distribution of material around the source, a tight upper limit on the column density $N_H < 10^{24} \text{cm}^{-2}$ derives from the requirement that the Thomson optical depth be less than a few, to avoid smearing out the the short time-scale structure of the GRB by Thomson scattering (Boettcher *et al.* 1999). Furthermore, Fe absorption features of strength comparable to that in emission should also be detected.

It is also conceivable that the impact of the relativistic shell that produced the original GRB on these ejecta could provide an additional energy input in the afterglow.

GRB970508 is the most promising candidate of the BeppoSAX afterglows for the search of line emission. It is characterized by a large outbursting event during its afterglow phase (Piro *et al.* 1998) and has the highest signal to noise ratio of the BeppoSAX GRB afterglows. In this Letter we present results of a spectral analysis of the afterglow of GRB970508 and discuss their implications on the nature of the central engine of GRB.

2. Observations of the X-ray afterglow spectrum of GB970508

The Gamma-ray burst GRB970508 was detected by the Gamma-ray Burst Monitor (GRBM) aboard BeppoSAX (Piro *et al.* 1998). The Wide Field Camera (WFC) provided a few arcmin position acquired with the Narrow Field Instruments 6 hours after the burst, which led to the identification of the X-ray afterglow associated with the GRB. Three further observations were carried out, the last performed 6 days after the burst (Fig.1, Tab.1).

LECS(0.1-3.5 keV) and MECS(1.8-10.0 keV) spectra of the observations were obtained using standard procedures (BeppoSAX web pages: www.sdc.asi.it), with 15-20 counts (source + background) per bin.

Inspection of the light curve (Fig.1) shows that in the first half period of observation number 1 the flux decays, followed by a rising trend in the second half. We have thus split this observation into two parts (1a & 1b) at the time of the minimum of the light curve, to investigate spectral evolution in different states, producing 5 spectra in total.

2.1. The iron K_α line

We fitted each spectrum with a power law with variable absorption, keeping the relative normalization of LECS vs MECS free to vary in the range 0.6-0.9. This model provides a good fit to all but dataset 1a. In this case we have $\chi^2_\nu = 2.4$ for $\nu = 8$ degrees of freedom, which corresponds to a probability of 1.4%¹

¹Note that GB970508 has the highest S/N ratio of the BeppoSAX GRB afterglows, so that the number of spectra with comparable statistical weight is limited to those of data set 1. We also remind that the afterglow of GB970228 had an higher flux, but the S/N ratio

However the major contribution to χ^2 is not randomly distributed but is concentrated in a line feature around 3.5 keV (Fig. 2a; note that in figure we show for clarity only data points below 2 KeV for LECS and above 2 keV for MECS).

The possible association of this feature with Fe K line emission is obvious, considering that its energy coincides with Fe I-Fe XX $K\alpha$ emission at the redshift $z=0.835$, i.e. that of the putative host galaxy (Bloom et al. 1998). We have therefore added to the power law model a narrow line (width $\leq 1.5 \times \text{FWHM} \simeq 0.5$ keV) at the energy of 3.5 keV. The relative normalization LECS/MECS – that depends primarily on the position of the source in the LECS detector has, in all datasets, an error greater than the nominal range 0.6-0.9. We have hereafter fixed it to 0.8, the value averaged over all data sets. The best fit now yields $\chi^2_\nu = 0.93$ with $\nu = 7$. The improvement of $\Delta\chi^2/\chi^2_\nu = 13.6$ corresponds to a confidence level of 99.3% (F test). We stress that this confidence level relies on the a priori knowledge of the energy of the Fe line available from an independent redshift measurement. Were it not available, the confidence level for the presence of a line with free energy would have been 97%.

The best fit line intensity is $I_{Fe} = (5 \pm 2) \times 10^{-5}$ ph cm $^{-2}$ s $^{-1}$ (hereafter all errors and upper limits correspond to 90% confidence level for a single parameter). Upper limits on line intensity in datasets 1b (Fig.2b), 2 3 and 4 are 2.5, 2, 2.5 and 0.5×10^{-5} ph cm $^{-2}$ s $^{-1}$ respectively.

The power law parameters obtained in dataset 1a and 1b are consistent with each other (see Tab. 1) and the absorption column densities are consistent with the hydrogen column in the line of sight of our Galaxy, i.e. the intrinsic absorption is consistent with zero. We next performed a simultaneous fit to both datasets with the same continuum parameters,

was lower, because data of one MECS only were usable (Costa *et al.* 1997).

and the addition of a line to dataset 1a only. In this case we left the energy of the line as free parameter, deriving $E = 3.4 \pm 0.3$, corresponding to (6.2 ± 0.6) in the source rest frame. The error range is too large to identify the ionization state of iron. In fig.2 (inset) we show the contour plot of I_{Fe} vs E .

We note that the absence of the line in dataset 1b explains why a power law model provides an acceptable fit to the summed spectrum (Owens *et al.* 1998). We have also investigated whether the spectral continuum could be fitted by a black body or a thermal bremsstrahlung. We find that thermal models provide worse results than that of a power law continuum, as already shown in the case of GRB970228 (Frontera *et al.* 1998).

2.2. Spectral variability in the afterglow outburst

Spectral variability of the afterglow spectra supports the idea that a refreshed energy input produced the outburst observed in the light curve. The spectral index in the first observation (Tab. 1) is $\alpha = 1.5 \pm 0.6$. In the second and third observation the flux lies above an extrapolation of the light curve expected from a power law evolution from the primary event. The spectrum is harder ($\alpha = 0.4 \pm 0.6$), indicating electron reacceleration, as expected in a newly formed shock. In the last observation the flux decreases and the spectrum is very steep ($\alpha = 2.2 \pm 0.7$): the new energy source has been exhausted. We have noticed previously that spectra 1a and 1b are consistent with each other. Since the transition 1a-1b identifies the beginning of the outburst, one would then expect an hardening of the spectrum in 1b. However such an effect could be masked by the continuation of the power law decay law (Fig.1), which accounts for about 60% of the flux observed in dataset 1b.

3. Origin of the line and constraints on the emitting region

The minimum amount of mass M_{min} present can be estimated (Lazzati *et al.* 1998) simply by counting how many ionizing photons a given iron atom can absorb, assuming that recombination is much faster than the time interval between successive ionizations, and by demanding that the existing iron atoms thus account for the fluence in the line:

$$M_{min} = 0.1 M_{\odot} A_{Fe}^{-1} \left(\frac{I_{Fe}}{10^{-5} \text{ ph s}^{-1} \text{ cm}^{-2}} \right) \frac{T}{10^5 \text{ s}} \left(\frac{R}{10^{16} \text{ cm}} \right)^2 \frac{0.1}{q} E_{52}^{-1} \quad (1)$$

where A_{Fe} is the iron abundance in units of the solar value, T is the line duration, $R = 10^{16} R_{16} \text{ cm}$ is the distance of the medium from the central source, and $q < 0.1$ is the fraction of the total energy $E = 10^{52} E_{52} \text{ erg}$ which is recycled into the line. Since the line disappears after about 10^5 s , we expect the emitting region to have a size $D \approx 3 \times 10^{15} \text{ cm}$; in fact, since the line width is mostly instrumental, we can bar the existence of relativistic effects. Then the typical density is $n = 5 \times 10^9 A_{Fe}^{-1} R_{16}^2 \text{ cm}^{-3}$, and typical Thompson and iron optical depths are $\tau_T = 10 A_{Fe}^{-1} R_{16}^2$ and $\tau_{Fe} = 5 R_{16}^2$.

From the above we deduce that the emitting material cannot lie along the line of sight: for $R_{16} > 1$, $\tau_T \gg 1$, and Thomson scattering would smear out the time structure of the burst, contrary to observations (Boettcher *et al.* 1999). Even if $A_{Fe} \gg 1$, we would still have $\tau_{Fe} \gg 1$, which implies an iron edge not seen in the spectrum. For $R_{16} < 1$, we notice that the fireball would load itself with so many baryons to spoil its relativistic expansion, again contradicting observations.

Having established that the emitting medium lies sideways from the line of sight, we also notice that we cannot have $R \gg D$, because then the line would appear much later than 1 day. We conclude that $R \approx D$.

The ionization parameter, using the minimum density derived above, is found to be

$$\xi = \frac{L}{n R^2} = 10^8 A_{Fe} \frac{L}{10^{50} \text{ erg s}^{-1}} R_{16}^{-4} \approx 10^9 A_{Fe} t^{-1.1} R_{16}^{-4} \quad (2)$$

where L is the luminosity in the range 0.016-10 keV, whose temporal evolution is approximately described by a power law (Piro *et al.* 1998). From Eq. (2) one sees that for early times ($t \lesssim 10^4$ s), the material will be fully ionized, and thus the line may be due to recombination, while at later times, when $\xi \lesssim 10^4$, fluorescence may account for the emission (Kallmann & McCray 1982, Hirano *et al.* 1987). The exact moment of the transition cannot be determined, because we established above that the emitting material lies sideways with respect to the observer, and we cannot exclude some amount of beaming, but the following discussion still stands. At early times, the emitting material will probably reach close to the Inverse-Compton temperature in the burst radiation field, which clearly exceeds 10^9 °K; for these high temperatures and low densities, recombination is not an effective emitter of line photons, and it is most likely that most of the line we see is emitted by fluorescence in the time range 10^4 s $< t < 10^5$ s, during which the total observed fluence is $\approx 10^{51}$ erg. Putting this into Eq. 1, we find a minimum mass of

$$M_{min} = 0.5 M_{\odot} A_{Fe}^{-1} \left(\frac{R_{16}}{0.3} \right)^2, \quad (3)$$

which is our best estimate for the total mass present.

3.1. The site of the line emission

In principle, the line might arise from the normal interstellar medium surrounding the burst volume, or from a normal stellar wind emitted by the burst progenitor. However, a very large density ($n \approx 10^9$ cm $^{-3}$) is necessary to account for the observed spectral feature; this is obvious when one considers that the line reported here is as bright as a typical one from clusters of galaxies at $z \approx 0.1 \ll z_{970508}$. Winds of massive stars are also unlikely to be responsible for the iron line: even for rather extreme wind parameters ($\dot{M} = \dot{m}_{-4} \times 10^{-4} M_{\odot}$ yr $^{-1}$, and $v = v_2 \times 100$ km s $^{-1}$), the wind density exceeds the minimum density only for $R_{16} < 0.1(\dot{m}_{-4} A_{Fe}/v_2)^{1/4}$. However, for this small distance, the ionization

parameter (Eq. 2) is $\gtrsim 10^7$ even a day after the burst and, even including beaming, the wind material will be totally ionized; under these conditions, only recombination can provide the line photons, but the recombination rates are very small ($t_{rec} > 10^5$ s), and thus the conditions of instantaneous de-excitation under which Eq. 1 was established are no longer satisfied: much larger masses are required than can be accounted for by the winds. Furthermore, such small distances ($R_{16} < 0.1$) cannot explain the line duration. We thus conclude that the material surrounding the burst must have been pre-ejected.

$A_{Fe} \approx 1$ is likely to be close to an upper limit to the iron abundance of the ISM or of a stellar wind at $z \simeq 1$. Scenarios concerning mergers of binaries then begin to appear unlikely, regardless of whether the binary is made of two neutron stars, a black hole and a neutron star, or a black hole and a white dwarf, because it has been established (Mészáros and Rees 1992) that they cannot pre-eject, even under the rosier assumptions, more than $\approx 10^{-4} M_{\odot}$ due to their tidal interactions, about three orders of magnitude less than the limit in Eq. 1. We remark that the SupraNova (Vietri & Stella 1998) provides the most appealing scenario, because of the large amount of iron material ejected by the supernova explosion preceding the burst. In fact, theoretical computations of the explosion of massive stars indicate that $\approx 0.1 M_{\odot}$ of iron are ejected by the SN (Woosley & Weaver 1995, Thielemann, Nomoto & Hashimoto 1996). In the case of SN1987a, whose progenitor had $M \approx 20 M_{\odot}$, a similar amount of iron was derived from optical observations (Danziger & Bouchet 1993).

In future, it is conceivable that simultaneous observations of the Ly- α line may help to determine A_{Fe} , and thus uniquely determine the total illuminated mass, and by implication the adequacy of different formation scenarios.

4. Summary

In this paper we have obtained the following results:

- We have searched the X-ray spectrum of GB970508’s afterglow for an iron line, located at the system’s redshift ($z = 0.835$); we found such a line with limited statistical significance (99.3%) in the early part (first 16 h) of the afterglow; the line decreases in the later part of the observations (≈ 1 day after the burst) by at least a factor 2, enough to make it undetectable with current apparatus;
- Simultaneously with the line disappearance, the X-ray flux both rises and hardens ($\alpha = 0.4 \pm 0.6$, while $\alpha = 1.5 \pm 0.6$ before the reburst), consistent with the appearance of a new shock. Then, at the end of the outburst, the spectrum steepens.

We showed that these observations indicate that a mass $\approx 0.5M_{\odot}A_{Fe}^{-1}$ should be located at a distance of 3×10^{15} cm and sideways with respect to the observer, and is moving at subrelativistic speed. In order to reach such distance, this material must have been pre-ejected by the source originating the burst, shortly (perhaps a year, for typical SN expansion speeds) before the burst. We stress that these observations contain two coincidences: on the one hand, this is the only burst in which a reburst and a line have been observed by BeppoSAX; on the other, the iron line disappears exactly at the moment of the reburst. We also point out that a line feature, with a similar significance, has been found by ASCA in another burst, GRB970828, which also shows an event of rebursting during the X-ray afterglow (Yoshida *et al.* 1998).

We hope the report of this result will help shape future observations by X-ray satellites with superior observing capabilities, such as XMM, AXAF, ASTRO-E.

We thank the referee, Dr. M. Boettcher, for his critical suggestions that improved the paper substantially, F. Matteucci for helpful discussions and the BeppoSAX team for the support with observations. BeppoSAX is a program of the Italian space agency (ASI) with the participation of the Dutch space agency (NIVR).

REFERENCES

- Bloom J.S., Djorgovski S.G., Kulkarni S. R. & Frail D. A. 1998, ApJ, **507**, 25.
- Boella G. *et al.* 1997, *A&AS*, **122**, 299.
- Boettcher M., Dermer C.D.D., Crider A. W. & Liang E. D. 1999, A&A, in press, preprint: astro-ph/9809156.
- Costa E. *et al.* 1997, Nature, **387**, 783
- Danziger J. & Bouchet, P. 1993, in New Aspects of Magellanic Cloud research, proc. of *Evolution of Galaxies*, Heidelberg, June 15-17 1992, p.208
- Frontera F. *et al.* 1998, ApJ, **493**, 67
- Galama T.J. *et al.* 1998 Nature, **395**, 670.
- Ghisellini G. *et al.* , ApJ, submitted; preprint: astro-ph/9808156.
- Hirano T. *et al.* 1987 PASJ, 39, 619
- Kallmann T. R. & McCray R. 1982, ApJS, 50 263
- Kulkarni S. R. *et al.* 1998, Nature, **395**, 663.
- Lazzati D., Ghisellini G. & Campana S., ApJ, submitted.
- Mészáros P. & Rees M. J. 1992, ApJ, **397**, 570.
- Mészáros P. & Rees M. J. 1998, MNRAS, **299**, L10.
- Metzger M.R. *et al.* 1997 Nature, **387**, 878.
- Owens A. *et al.* 1998, A&A, **339**, L37.

- Paczynski, B. 1998, ApJ, **494**, L45.
- Perna R. & Loeb A., 1998, ApJ, **501**, 467.
- Piro L., Scarsi L. & Butler R.C. in *X-Ray and EUV/FUV Spectroscopy and Polarimetry* (ed. S. Fineschi) SPIE 2517, 169-181 (1995).
- Piro L. *et al.* A&A1998 **331**, L41.
- Thieleman F-K., Nomoto, K. & Hashimoto, M. 1996, ApJ, **460**, 408
- van Paradijs *et al.* 1997, Nature, **386**, 686.
- Vietri M. & Stella L. ApJ, **507**, L45 (1998)
- Woosley S. E.& Weaver, T. A. 1995 ApJS, **101**, 181.
- Yoshida A. *et al.* 1998, Proc. of *Gamma-ray bursts in the afterglow era*, Rome, Nov. 3-6 1998, Frontera F. & Piro L. eds., A&AS, submitted

Fig. 1.— The X-ray (2-10 keV) light curve of the X-ray afterglow of GRB980508. The dashed line represents the extrapolation from the X-ray data points of the burst (Piro *et al.* 1998). For the spectral analysis data from the first pointing was divided in two parts, corresponding to the transition between the decay and turn up from the light curve, creating in total 5 spectra (1a,1b-4).

Fig. 2.— The spectra (in detector counts) of the afterglow of GRB970508 from data set 1a (panel a) and data set 1b (panel b). The continuous line represents the best fit power law continuum in the range 0.1-10 keV, excluding the interval 3-4 keV. For clarity we show in figure LECS data below 2 keV and MECS data above 2 keV. The inset in panel a) shows the contour plot of the line intensity vs energy. Contours correspond to 68%, 90% & 99% confidence levels for two interesting parameters.

Table 1. Log of Observations and fit with a power law

Obs.	T_{start}^a	Exp.Time (ksec)		Cnt Rate (10^{-3} cts s $^{-1}$)		α	N_H (10^{22} cm $^{-2}$)	χ^2/ν
		LECS	MECS	LECS(0.1-2 keV)	MECS(2-10 keV)			
1a ^b	0.24	8	11	3.0 ± 0.8	8.8 ± 1.2	1.5 ± 0.9	$0.6^{+1.3}_{-0.55}$	6.5/7
1b	0.65	7	16	1.4 ± 0.7	7.2 ± 0.9	1.7 ± 1.0	$1.3^{+2.5}_{-1.25}$	6/9
1a+b ^b	0.24	15	28	2.1 ± 0.5	8.3 ± 0.8	1.5 ± 0.6	$0.7^{+1.5}_{-0.65}$	14.2/18
2	2.74	6	24	1.3 ± 0.7	5.7 ± 0.7	0.4 ± 0.6	$0.05^{+1.5}$	6.2/9
3	4.14	3	12	< 1.5	3.1 ± 0.9	0.5 ± 0.8	0.05 <i>fix</i>	5.2/6
4	5.7	14	73	1.7 ± 0.5	1.2 ± 0.3	2.2 ± 0.7	$0.05^{+0.5}$	6.3/10

Errors of spectral parameters correspond to 90% confidence level for a single parameter of interest; $N_{HGal} = 5 \cdot 10^{20} \text{cm}^{-2}$

^astart of the observation in days from the GRB

^bincluding a line in spectral fitting of dataset 1a



